arXiv:1111.4849v1 [math.NT] 21 Nov 2011

THE PROPERTIES OF MODIFIED q-BERNSTEIN POLYNOMIALS FOR FUNCTIONS OF SEVERAL VARIABLES WITH THEIR GENERATING FUNCTION AND INTERPOLATION FUNCTION

Mehmet AÇIKGÖZ[†], **Serkan ARACI**[†] **and Hassan JOLANY**[‡]

[†]University of Gaziantep, Faculty of Arts and Science, Department of Mathematics, 27310
Gaziantep, Turkey
[‡]School of Mathematics, Statistics and Computer Science, University of Tehran, Iran
acikgoz@gantep.edu.tr; mtsrkn@hotmail.com; hassan.jolany@khayam.ut.ac.ir

Abstract

The aim of this paper is to give a new approach to modified q-Bernstein polynomials for functions of several variables. By using these polynomials, recurrence formulas and some new interesting identities related to the second Stirling numbers and generalized Bernoulli polynomials are derived. Moreover, the generating function and interpolation function of these polynomials of several variables and also the derivatives of these polynomials and their generating function are given.

2000 Mathematics Subject Classification 11M06, 11B68, 11S40, 11S80, 28B99, 41A50

Key Words and Phrases Generating function, Bernstein polynomial of several variables, Bernstein operator of several variables, shift difference operator, q-difference operator, second kind Stirling numbers, generalized Bernoulli polynomials, Mellin transformation, interpolation function

1. Introduction, Definitions and Notations

In approximation theory, the Bernstein polynomials, named after their creater S. N. Bernstein in 1912, have been studied by many researchers for a long time. But no one knew anything about the generating function of these polynomials until recently Simsek and Acikgoz constructed a generating function of (q-) Bernstein type polynomials based on the q-analysis, [14], and gave some new relations related to these polynomials, Hermite polynomials, Bernoulli polynomials of higher order and the second kind Stirling numbers. Interpolation function of (q-) Bernstein type polynomials is defined by applying Mellin transformation to this generating function. They gave some relations and identities on these polynomials. Later then, the Acikgoz and Araci constructed the generating function for classical Bernstein polynomials, and for Bernstein polynomials for functions of two variables and gave their properties (see [1], [2], [3], and [15] for details). Recently, T. Kim, J. Choi and Y. H. Kim have studied on the k-dimensional generalization of q-Bernstein polynomials, in which they have given some interesting properties of the k-dimensional generalization of q-Bernstein polynomials (for details, see[8]). Our generalization of q-Bernstein polynomials are different from the k-dimensional generalization of q-Bernstein polynomials of T. Kim, J. Choi and Y. H. Kim. In the present paper, we also derived some interesting properties of our generalization of q-Bernstein polynomials.

Now, we are ready to give some definitions and some properties of Bernstein polynomials of several variables with their generating function.

Let $C\left(\underbrace{D\times D\times ...\times D}_{w-times}\right)$ denotes the set of continuous functions on $\underbrace{D\times D\times ...\times D}_{w-times}=\underbrace{[0,1]\times[0,1]\times ...\times[0,1]}_{w-times}$. For $f\in C\left(\underbrace{D\times D\times ...\times D}_{w-times}\right)$, we have

$$\mathbf{B}_{n_1,n_2,\cdots,n_w}\left(f;x_1,x_2,\cdots,x_w\right) \; : \; =\sum_{k_1=0}^{n_1}\sum_{k_2=0}^{n_2}\cdots\sum_{k_W=0}^{n_W} f\left(\frac{k_1}{n_1},\frac{k_2}{n_2},\cdots,\frac{k_w}{n_w}\right) \times B_{k_1,k_2,\cdots,k_w;n_1,n_2,\cdots,n_w}\left(x_1,x_2,\cdots,x_w\right) \tag{1.1}$$

where $\mathbf{B}_{n_1,n_2,\cdots,n_w}\left(f;x_1,x_2,\cdots,x_w\right)$ is called the Bernstein operator of several variables of order $\sum\limits_{i=1}^{w} n_i$ for f. For $k_i,n_i\in\mathbb{N}_0$ with $i=1,2,..w$, the Bernstein polynomials of several variables of degree $\sum\limits_{i=1}^{w} n_i$ is defined by

$$B_{k_1,k_2,\cdots,k_w;n_1,n_2,\cdots,n_w}\left(x_1,x_2,\cdots,x_w\right)=\prod_{i=1}^{w}\binom{n_i}{k_i}x_i^{k_i}\left(1-x_i\right)^{n_i-k_i}, \tag{1.2}$$

where $\binom{n}{k}=\frac{n(n-1)\cdots(n-k+1)}{k!}$ and $x_i\in D$ for $i=1,2,...,w$. These polynomials satisfy the relation

$$B_{k_1,k_2,\cdots,k_w;n_1,n_2,\cdots,n_w}\left(x_1,x_2,\cdots,x_w\right)=\prod_{i=1}^{w}B_{k_i,n_i}\left(x_i\right)$$

and they have form a partition of unity; that is

$$\sum_{k_1=0}^{n_1}\sum_{k_2=0}^{n_2}\cdots\sum_{k_w=0}^{n_w}B_{k_1,k_2,\cdots,k_w;n_1,n_2,\cdots,n_w}\left(x_1,x_2,\cdots,x_w\right)=1.$$

By using the definition of Bernstein polynomials for functions of several variables, it is not difficult to prove the property given above as

$$\sum_{k_1=0}^{n_1}\sum_{k_2=0}^{n_2}\cdots\sum_{k_w=0}^{n_w}\prod_{i=1}^{w}B_{k_i,n_i}\left(x_i\right)=1.$$

Also, $B_{k_1,k_2,\cdots,k_w;n_1,n_2,\cdots,n_w}\left(x_1,x_2,\cdots,x_w\right)=0$ for $k_i>n_i$ with $i=1,2,...,w$, because $\binom{n_i}{k_i}=0$. There are $\prod\limits_{i=1}^{w}\left(n_i+1\right)$, $\sum\limits_{i=1}^{w} n_i$-th degree Bernstein polynomials.

Many researchers have studied the Bernstein polynomials of two variables in approximation theory (see [5], [6]). But nothing was known about the generating function of these polynomials. Note that for $k_i, n_i \in \mathbb{N}_0$ and $x_i \in D$ with $i = 1, 2, ..., w$, we obtain the generating function for $B_{k_1,k_2,\cdots,k_w;n_1,n_2,\cdots,n_w}(x_1, x_2, \cdots, x_w)$ as follows:

$$\begin{aligned} F_{k_1,k_2,\cdots,k_w}(t; x_1, x_2, \cdots, x_w) &= \prod_{i=1}^{w} \frac{(tx_i)^{k_i}}{k_i!} e^{wt - t\sum\limits_{i=1}^{w} x_i} \qquad (1.3) \\ &= \sum_{n_1=k_1}^{\infty} \sum_{n_2=k_2}^{\infty} \cdots \sum_{n_w=k_w}^{\infty} B_{k_1,k_2,\cdots,k_w;n_1,n_2,\cdots,n_w}(x_1, x_2, \cdots, x_w) \prod_{i=1}^{w} \frac{t^{n_i}}{n_i!} \end{aligned}$$

where

$$B_{k_1,k_2,\cdots,k_w;n_1,n_2,\cdots,n_w}(x_1, x_2, \cdots, x_w) = \begin{cases} \prod\limits_{i=1}^{w} \binom{n_i}{k_i} x_i^{k_i} (1 - x_i)^{n_i - k_i} & , \text{ if } \ n_i \geq k_i \\ 0 & , \text{ if } \ n_i < k_i \end{cases}$$

for $k_i \in \mathbb{N}_0$ and $x_i \in D$, for, $i = 1, 2, ..., w$.

Remark 1. *By substituting* $w = 1$ *into (1.3), we get a special case of* $F_{k_1,k_2,\cdots,k_w}(t; x_1, x_2, \cdots, x_w)$ *which was proved by Acikgoz and Araci (for details, see* [1]*)*

$$F_{k_1}(t, x_1) = \frac{(tx_1)^{k_1} e^t}{k_1! e^{tx_1}} = \sum_{n_1=k_1}^{\infty} B_{k_1,n_1}(x_1) \frac{t^{n_1}}{n_1!}.$$

Let $0 < q < 1$. Define the q-number of x by $[x]_q := \frac{1-q^x}{1-q}$ and $[x]_{-q} := \frac{1-(-q)^x}{1+q}$, (see [4], [7], [8], [9], [10], [12], [13], [14], [15], for details and related facts). Note that $\lim\limits_{q\to 1}[x]_q = x$. [7] is actually motivated the authors to write this paper and they have extended all results given in [7] to modified q-Bernstein polynomials of several variables.

2. The Modified q-Bernstein Polynomials for Functions of Several Variables

For $0 < q < 1$, consider

$$\begin{aligned} F_{k_1,k_2,\cdots,k_w}(t, q; x_1, x_2, \cdots, x_w) &= \prod_{i=1}^{w} \frac{\left(t [x]_q\right)^{k_i}}{k_i!} e^{t\sum\limits_{i=1}^{w}[1-x_i]} \\ &= \sum_{n_1=k_1}^{\infty} \sum_{n_2=k_2}^{\infty} \cdots \sum_{n_w=k_w}^{\infty} B_{k_1,k_2,\cdots,k_w;n_1,n_2,\cdots,n_w}(x_1, x_2, \cdots, x_w; q) \prod_{i=1}^{w} \frac{t^{n_i}}{n_i!} \end{aligned}$$

where $k_i, n_i \in \mathbb{N}_0$ and $x_i \in D$ for $i = 1, 2, ..., w$. We note that

$$\lim_{q\to 1} F_{k_1,k_2,\cdots,k_w}(t, q; x_1, x_2, ..., x_w) = F_{k_1,k_2,\cdots,k_w}(t; x_1, x_2, ..., x_w).$$

Definition 1. *We define the generating function of modified q-Bernstein polynomials for functions of several variables as follows:*

$$\begin{aligned} F_{k_1,k_2,\cdots,k_w}(t,q;x_1,x_2,\cdots,x_w) &= \prod_{i=1}^{w}\frac{\left(t\,[x]_q\right)^{k_i}}{k_i!}e^{t\sum\limits_{i=1}^{w}[1-x_i]} \qquad (2.1)\\ &= \sum_{n_1=k_1}^{\infty}\sum_{n_2=k_2}^{\infty}\cdots\sum_{n_w=k_w}^{\infty}B_{k_1,k_2,\cdots,k_w;n_1,n_2,\cdots,n_w}(x_1,x_2,\cdots,x_w;q)\prod_{i=1}^{w}\frac{t^{n_i}}{n_i!} \end{aligned}$$

where $k_i, n_i \in \mathbb{N}_0$ and $x_i \in D$ with $i = 1, 2, ..., w$.

By using the taylor expansion of $e^{t\sum\limits_{i=1}^{w}[1-x_i]}$ and the comparing coefficients on the both sides in (2.1), we simply get the following Corollary 1:

Corollary 1. *For $k_i, n_i \in \mathbb{N}_0$ and $x_i \in D$ for $i = 1, 2, ..., w$;*

$$\begin{aligned} &B_{k_1,k_2,\cdots,k_w;n_1,n_2,\cdots,n_w}(x_1,x_2,\cdots,x_w;q)\\ &= \begin{cases} \prod\limits_{i=1}^{w}\binom{n_i}{k_i}[x_i]_q^{k_i}[1-x_i]_q^{n_i-k_i} & , \text{ if } n_i \geq k_i \text{ for } i\in[1,w] \\ 0 & , \text{ if } n_i < k_i \text{ for } i\in[1,w] \end{cases}. \end{aligned} \tag{2.2}$$

Theorem 1. ***(Recurrence Formula for** $B_{k_1,k_2,\cdots,k_w;n_1,n_2,\cdots,n_w}(x_1,x_2,\cdots,x_w;q)$**)** For $k_i, n_i \in \mathbb{N}_0$, $x_i \in D$, and $i = 1, 2, ..., w$, we have*

$$B_{k_1,k_2,\cdots,k_w;n_1,n_2,\cdots,n_w}(x_1,x_2,\cdots,x_w;q) = \prod_{i=1}^{w}[1-x_i]B_{k_i;n_i-1}(x_i;q)+[x_i]B_{k_i-1;n_i-1}(x_i;q). \tag{2.3}$$

Proof. By using the definition of Bernstein polynomials for functions of several variables, we have

$$\begin{aligned} B_{k_1,k_2,\cdots,k_w;n_1,n_2,\cdots,n_w}(x_1,x_2,\cdots,x_w;q) &= \prod_{i=1}^{w}\left[\binom{n_i-1}{k_i}+\binom{n_i-1}{k_i-1}\right][x_i]_q^{k_i}[1-x_i]_q^{n_i-k_i}\\ &= \prod_{i=1}^{w}[1-x_i]B_{k_i;n_i-1}(x_i;q)+[x_i]B_{k_i-1;n_i-1}(x_i;q). \end{aligned}$$

This is the desired result. ☐

Remark 2. *By setting $w = 1$ and $q \to 1$ into (2.3), we get the familiar identity $B_{k_1,n_1}(x_1)$ for as follows:*

$$B_{k_1,n_1}(x_1) = (1-x_1)\,B_{k_1,n_1}(x_1)+x_1 B_{k_1,n_1}(x_1).$$

(see [1], [3], [14]*).*

Theorem 2. *For $k_i, n_i \in \mathbb{N}_0$ and $x_i \in D$ for $i = 1, 2, ..., w$, we have*

$$\begin{aligned} & B_{n_1-k_1,n_2-k_2\cdots,n_w-k_w;n_1,n_2,\cdots,n_w}\left(1-x_1,1-x_2,\cdots,1-x_w;q\right) \\ = \; & B_{k_1,k_2,\cdots,k_w;n_1,n_2,\cdots,n_w}\left(x_1,x_2,\cdots,x_w;q\right) \qquad (2.4) \end{aligned}$$

Remark 3. *By substituting $w = 1$ into (2.4), we get the well-known identity as follows:*

$$B_{n_1-k_1,n_1}\left(1-x_1\right) = B_{k_1,n_1}\left(x_1\right).$$

(see [1], [3]*).*

Definition 2. *Let f be a continuous function of several variables on $\underbrace{D \times D \times ... \times D}_{w-times}$. Then the modified q-Bernstein operator of order $\sum_{i=1}^{w} n_i$ for f is defined by*

$$\begin{aligned} & \mathbf{B}_{n_1,n_2,\cdots,n_w}\left(f : x_1,x_2,\cdots,x_w;q\right) \qquad (2.5) \\ = \; & \sum_{k_1=0}^{n_1}\sum_{k_2=0}^{n_2}\cdots\sum_{k_W=0}^{n_W} f\left(\frac{k_1}{n_1},\frac{k_2}{n_2},\cdots,\frac{k_w}{n_w}\right) B_{k_1,k_2,\cdots,k_w;n_1,n_2,\cdots,n_w}\left(x_1,x_2,\cdots,x_w;q\right) \end{aligned}$$

where $x_i \in D$, $n_i \in \mathbb{N}$.

When we set $f\left(\frac{k_1}{n_1},\frac{k_2}{n_2},\cdots,\frac{k_w}{n_w}\right) = 1$ into (2.5), we easily see that,

$$\begin{aligned} & \mathbf{B}_{n_1,n_2,\cdots,n_w}\left(1 : x_1,x_2,\cdots,x_w;q\right) \qquad (2.6) \\ = \; & \sum_{k_1=0}^{n_1}\sum_{k_2=0}^{n_2}\cdots\sum_{k_W=0}^{n_W} B_{k_1,k_2,\cdots,k_w;n_1,n_2,\cdots,n_w}\left(x_1,x_2,\cdots,x_w;q\right) \end{aligned}$$

From the definition binomial theorem and (2.6), we get the following Corollary 2 for modified q-Bernstein polynomials for functions of several variables:

Corollary 2. *For any $k_i, n_i \in \mathbb{N}_0$ and $x_i \in D$ for $i = 1, 2, ..., w$, we have*

$$\mathbf{B}_{n_1,n_2,\cdots,n_w}\left(1 : x_1,x_2,\cdots,x_w;q\right) = \prod_{i=1}^{w}\left(1+(1-q)\,[x_i]_q[1-x_i]_q\right)^{n_i}, \qquad (2.7)$$

we easily see for $\lim_{q\to 1}\mathbf{B}_{n_1,n_2,\cdots,n_w}\left(1 : x_1,x_2,\cdots,x_w;\right) = 1$. This is partition of unity for modified Bernstein polynomials for functions of several variables.

Theorem 3. *For $\xi_j \in \mathbb{C}$, $x_j \in D$, and for $n_j \in \mathbb{N}$, with $j = 1, 2, ..., w$, we have*

$$B_{k_1,k_2,\cdots,k_w;n_1,n_2,\cdots,n_w}\left(x_1,x_2,\cdots,x_w;q\right) = \frac{1}{(2\pi i)^w}\underbrace{\oint_C\oint_C\cdots\oint_C}_{w\text{-}times}\prod_{j=1}^{w} n_j!F_q^{(k_j)}\left(x_j,\xi_j\right)\frac{d\xi_j}{\xi_j^{n_j+1}} \qquad (2.8)$$

where

$$F_q^{(k)}\left(x,t\right) = \frac{\left(t[x]_q\right)^k}{k!}e^{t[1-x]_q} \text{ (see [14], [15])}$$

and C is a circle around the origin and integration is in the positive direction.

Proof. By using the definition of the modified q-Bernstein polynomials of several variables and the basic theory of complex analysis including Laurent series that

$$\begin{aligned}
&\underbrace{\oint_C \oint_C \cdots \oint_C}_{w\text{-times}} \prod_{j=1}^{w} F_q^{(k_j)}\left(x_j,\xi_j\right)\frac{d\xi_j}{\xi_j^{n_j+1}} \\
&= \sum_{l_1=0}^{\infty}\sum_{l_2=0}^{\infty}\cdots\sum_{lw=0}^{\infty}\oint_C\oint_C\cdots\oint_C\prod_{j=1}^{w}\frac{B_{k_j,l_j}\left(x_j,q\right)\xi_j^{l_j}}{l!}\frac{d\xi_j}{\xi_j^{n_j+1}} \\
&= (2\pi i)^w\left(\frac{B_{k_1,k_2,\cdots,k_w;n_1,n_2,\cdots,n_w}\left(x_1,x_2,\cdots,x_w;q\right)}{n_1!n_2!\cdots n_w!}\right). \qquad (2.9)
\end{aligned}$$

By using (2.9), we obtain

$$\begin{aligned}
&B_{k_1,k_2,\cdots,k_w;n_1,n_2,\cdots,n_w}\left(x_1,x_2,\cdots,x_w;q\right) \\
&= \frac{1}{(2\pi i)^w}\underbrace{\oint_C \oint_C \cdots \oint_C}_{w\text{-times}} \prod_{j=1}^{w} n_j!F_q^{(k_j)}\left(x_j,\xi_j\right)\frac{d\xi_j}{\xi_j^{n_j+1}}
\end{aligned}$$

and

$$\underbrace{\oint_C \oint_C \cdots \oint_C}_{w\text{-times}} \prod_{j=1}^{w} F_q^{(k_j)}\left(x_j,\xi_j\right)\frac{d\xi_j}{\xi_j^{n_j+1}} = (2\pi i)^w\left(\prod_{j=1}^{w}\frac{[x_j]_q^{k_j}\left[1-x_j\right]_q^{n_j-k_j}}{k_j!\left(n_j-k_j\right)!}\right). \qquad (2.10)$$

We also obtain from (2.4) and (2.10) that

$$\frac{1}{(2\pi i)^w}\underbrace{\oint_C \oint_C \cdots \oint_C}_{w\text{-times}} \prod_{j=1}^{w} n_j!F_q^{(k_j)}\left(x_j,\xi_j\right)\frac{d\xi_j}{\xi_j^{n_j+1}} = \prod_{j=1}^{w}\binom{n_j}{k_j}[x_j]_q^{k_j}[1-x_j]_q^{n_j-k_j}. \qquad (2.11)$$

Therefore we see from (2.9) and (2.11) that

$$B_{k_1,k_2,\cdots,k_w;n_1,n_2,\cdots,n_w}\left(x_1,x_2,\cdots,x_w;q\right) = \prod_{j=1}^{w}\binom{n_j}{k_j}[x_j]_q^{k_j}[1-x_j]_q^{n_j-k_j}.$$

Thus, we complete the proof of theorem. □

Theorem 4. *(**The Derivative Formula for** $B_{k_1,k_2,\cdots,k_w;n_1,n_2,\cdots,n_w}\left(x_1,x_2,\cdots,x_w;q\right)$) For $k_i \in \mathbb{N}$, $x_i \in D$, and for $i = 1,2,...,w$, we have*

$$\begin{aligned}
&\frac{\partial^w}{\partial x_1\partial x_2\cdots\partial x_w}\left(B_{k_1,k_2,\cdots,k_w;n_1,n_2,\cdots,n_w}\left(x_1,x_2,\cdots,x_w;q\right)\right) \\
&= \frac{\partial^w}{\partial x_1\partial x_2\cdots\partial x_w}\left(\prod_{i=1}^{w}\binom{n_i}{k_i}[1-x_i]_q^{k_i}[x_i]_q^{n_i-k_i}\right).
\end{aligned}$$

Proof. Using the definition of modified q-Bernstein polynomials for functions of several variables, the proof follows. $\square$

Therefore, we can write the modified q-Bernstein polynomials for functions of several variables as a linear combination of polynomials of higher order as follows:

Theorem 5. *For $k_i, n_i \in \mathbb{N}_0$, $x_i \in D$, and for $i = 1, 2, ..., w$, we have*

$$\begin{aligned} & B_{k_1,k_2,\cdots,k_w;n_1,n_2,\cdots,n_w}(x_1, x_2, \cdots, x_w; q) \\ = & \prod_{i=1}^{w}\left[\left(\frac{n_i - k_i + 1}{k_i}\right)\frac{[x_i]_q}{[1-x_i]_q}\right] B_{k_1-1,k_2-1,\cdots,k_w-1;n_1,n_2,\cdots,n_w}(x_1, x_2, \cdots, x_w; q). \end{aligned}$$

Proof. Using the definition of modified q-Bernstein polynomials for functions of several variables and the property (2.2), the proof follows. $\square$

Theorem 6. *If $n_i, k_i \in \mathbb{N}_0$ and $x_i \in D$ with $i = 1, 2, ..., w$, we have*

$$\begin{aligned} & B_{k_1,k_2,\cdots,k_w;n_1,n_2,\cdots,n_w}(x_1, x_2, \cdots, x_w; q) \\ = & \sum_{l_1=k_1}^{n_1}\sum_{l_2=k_2}^{n_2}\cdots\sum_{l_w=k_w}^{n_w}\prod_{i=1}^{w}\binom{n_i}{l_i}\binom{l_i}{k_i}(-1)^{l_i-k_i}q^{(l_i-k_i)(1-x_i)}[x_i]_q^{l_i}. \end{aligned}$$

Proof. From the definition of modified q-Bernstein polynomials of several variables and binomial theorem with n_i, $k_i \in \mathbb{N}_0$ and $x_i \in D$ for $i = 1, 2, ..., w$, we have

$$\begin{aligned} B_{k_1,k_2,\cdots,k_w;n_1,n_2,\cdots,n_w}(x_1, x_2, \cdots, x_w; q) &= \prod_{i=1}^{w}\binom{n_i}{k_i}[1-x_i]_q^{k_i}[x_i]_q^{n_i-k_i} \\ &= \sum_{l_1=k_1}^{n_1}\sum_{l_2=k_2}^{n_2}\cdots\sum_{l_w=k_w}^{n_w}\prod_{i=1}^{w}\binom{n_i}{l_i}\binom{l_i}{k_i}(-1)^{l_i-k_i}q^{(l_i-k_i)(1-x_i)}[x_i]_q^{l_i}. \end{aligned}$$

This is the desired result. $\square$

Theorem 7. *For $n_i, l_i \in \mathbb{N}_0$ and $x_i \in D$, with $i = 1, 2, ..., w$, we have*

$$\begin{aligned} & \left(\prod_{i=1}^{w}[x_i]_q\right)^m \\ = & \prod_{i=1}^{w}\frac{1}{\left([1-x_i]_q + [x_i]_q\right)^{n_i-l_i}}\sum_{k_1=m-1}^{n_1}\sum_{k_2=m-1}^{n_2}\cdots\sum_{k_w=m-1}^{n_w}\prod_{i=1}^{w}\frac{\binom{k_i}{m}}{\binom{n_i}{m}}B_{k_1,k_2,\cdots,k_w;n_1,n_2,\cdots,n_w}(x_1, x_2, \cdots, x_w; q). \end{aligned}$$

Proof. We easily see from the property of the modified q-Bernstein polynomials of several variables that

$$\sum_{k_1=1}^{n_1}\sum_{k_2=1}^{n_2}\cdots\sum_{k_w=1}^{n_w}\prod_{i=1}^{w}\frac{k_i}{n_i}B_{k_1,k_2,\cdots,k_w;n_1,n_2,\cdots,n_w}(x_1, x_2, \cdots, x_w; q) = \prod_{i=1}^{w}[x_i]_q\left([x_i]_q + [1-x_i]_q\right)^{n_i-1}$$

and that

$$\begin{aligned}
&\sum_{k_1=1}^{n_1}\sum_{k_2=1}^{n_2}\cdots\sum_{k_w=1}^{n_w}\prod_{i=1}^{w}\frac{\binom{k_i}{2}}{\binom{n_i}{2}}B_{k_1,k_2,\cdots,k_w;n_1,n_2,\cdots,n_w}\left(x_1,x_2,\cdots,x_w;q\right)\\
=\;&\left(\prod_{i=1}^{w}[x_i]_q\right)^2\left([x_i]_q+[1-x_i]_q\right)^{n_i-2}.
\end{aligned}$$

Continuing this method, we have

$$\begin{aligned}
\left(\prod_{i=1}^{w}[x_i]_q\right)^m \;=\;& \prod_{i=1}^{w}\frac{1}{\left([1-x_i]_q+[x_i]_q\right)^{n_i-l_i}}\\
&\times\sum_{k_1=m-1}^{n_1}\sum_{k_2=m-1}^{n_2}\cdots\sum_{k_w=m-1}^{n_w}\prod_{i=1}^{w}\frac{\binom{k_i}{m}}{\binom{n_i}{m}}B_{k_1,k_2,\cdots,k_w;n_1,n_2,\cdots,n_w}\left(x_1,x_2,\cdots,x_w;q\right)
\end{aligned}$$

and after making some algebraic operations, we obtain the desired result. □

We have seen from the theorem given above, it is possible to write $\left(\prod_{i=1}^{w}[x_i]_q\right)^m$ as a linear combination of modified q-Bernstein polynomials of several variables by using the degree evaluation formulae and mathematical induction method.

For $k\in\mathbb{N}_0$, the Bernoulli polynomials of degree k are defined by

$$\left(\frac{t}{e^t-1}\right)^k e^{xt}=\underbrace{\left(\frac{t}{e^t-1}\right)\left(\frac{t}{e^t-1}\right)\times\cdots\times\left(\frac{t}{e^t-1}\right)}_{k-times}e^{xt}=\sum_{n=0}^{\infty}B_n^{(k)}\left(x\right)\frac{t^n}{n!},$$

and $B_n^{(k)}=B_n^{(k)}\left(0\right)$ are called the n-th Bernoulli numbers of order k. It is well known that the second kind Stirling numbers are defined by $\frac{\left(e^t-1\right)^k}{k!}:=\sum_{n=0}^{\infty}S\left(n,k\right)\frac{t^n}{n!}$ for $k\in\mathbb{N}$ (see [7] and [14]). By using the above relations we can give the following theorem:

Theorem 8. *For $k_i,n_i,\in\mathbb{N}_0$ and $x_i\in D$ with $i=1,2,...,w$, we have*

$$\begin{aligned}
&B_{k_1,k_2,\cdots,k_w;n_1,n_2,\cdots,n_w}\left(x_1,x_2,\cdots,x_w;q\right)\\
=\;&\sum_{k_1=0}^{n_1}\sum_{k_2=0}^{n_2}\cdots\sum_{k_w=0}^{n_w}\prod_{i=1}^{w}[x_i]_q^{k_i}\binom{n_i}{l_i}B_{l_i}^{(k_i)}\left([1-x_i]_q\right)S\left(n_i-l_i,k_i\right).
\end{aligned}$$

Proof. By using the generating function of modified q-Bernstein polynomials of several variables, we have

$$\begin{aligned}\prod_{i=1}^{w}\frac{(t[x_i]_q)^{k_i}}{k_i!}e^{t\left(\sum_{i=1}^{w}[1-x_i]_q\right)} &= \prod_{i=1}^{w}[x_i]_q^{k_i}\left(\sum_{n_1=0}^{\infty}S(n_1,k_1)\frac{t^{n_1}}{n_1!}\right)\cdots\left(\sum_{n_w=0}^{\infty}S(n_w,k_w)\frac{t^{n_w}}{n_w!}\right)\times\\ &\quad\left(\sum_{l_1=0}^{\infty}B_{l_1}^{(k_1)}([1-x_1]_q)\frac{t^{l_1}}{l_1!}\right)\cdots\left(\sum_{l_w=0}^{\infty}B_{l_w}^{(k_w)}([1-x_w]_q)\frac{t^{l_w}}{l_w!}\right).\end{aligned}$$

By using the Cauchy product for sums given above

$$\begin{aligned}&B_{k_1,k_2,\cdots,k_w;n_1,n_2,\cdots,n_w}(x_1,x_2,\cdots,x_w;q)\\ &= \sum_{k_1=0}^{n_1}\sum_{k_2=0}^{n_2}\cdots\sum_{k_w=0}^{n_w}\prod_{i=1}^{w}[x_i]_q^{k_i}\binom{n_i}{l_i}B_{l_i}^{(k_i)}\left([1-x_i]_q\right)S(n_i-l_i,k_i).\end{aligned}$$

By comparing the last two relations, we have the desired result. □

Let Δ be the shift difference operator defined by $\Delta f(x)=f(x+1)-f(x)$. By using the mathematical induction method we have

$$\Delta^n f(0)=\sum_{k=0}^{n}\binom{n}{k}(-1)^{n-k}f(k), \tag{2.12}$$

for $n\in\mathbb{N}$ and using (2.12) in the generating function of second kind Stirling numbers,

$$\begin{aligned}\sum_{n=0}^{\infty}S(n,k)\frac{t^n}{n!} &= \frac{1}{k!}\sum_{l=0}^{k}\binom{k}{l}(-1)^{k-l}e^{lt}\\ &= \sum_{n=0}^{\infty}\frac{\Delta^k 0^n}{k!}\frac{t^n}{n!}. \qquad (2.13)\end{aligned}$$

By comparing the coefficients on both sides, we have

$$S(n,k)=\frac{\Delta^k 0^n}{k!}. \tag{2.14}$$

For $n_i,k_i\in\mathbb{N}$, by using the equation (2.14), we obtain the relation

$$\begin{aligned}&B_{k_1,k_2,\cdots,k_w;n_1,n_2,\cdots,n_w}(x_1,x_2,\cdots,x_w;q)\\ &= \sum_{k_1=0}^{n_1}\sum_{k_2=0}^{n_2}\cdots\sum_{k_w=0}^{n_w}\prod_{i=1}^{w}[x_j]_q^{k_j}\binom{n_i}{l_i}B_{l_i}^{(k_i)}\left([1-x_i]_q\right)\frac{\Delta^{k_i}0^{n_i}}{k_i!}\end{aligned}$$

which is the relation of the q-Bernstein polynomials of several variables in terms of Bernoulli polynomials of order k and second Stirling numbers with shift difference operator.

Let $(Eh)(x)=h(x+1)$ be the shift operator. Then the q-difference operator is defined by

$$\Delta_q^n=\prod_{i=0}^{n-1}\left(E-q^iI\right) \tag{2.15}$$

where I is the identity operator (see [7]).

For $f \in C[0,1]$ and $n \in \mathbb{N}$, we have

$$\Delta_q^n f(0) = \sum_{k=0}^{n} \binom{n}{k}_q (-1)^k q^{\binom{n}{2}} f(n-k), \tag{2.16}$$

where $\binom{n}{k}_q$ is the Gaussian binomial coefficient defined by

$$\binom{n}{k}_q = \frac{[n]_q [n-1]_q \cdots [n-k+1]_q}{[k]_q!}. \tag{2.17}$$

Theorem 9. *For $n_i, l_i \in \mathbb{N}_0$ and $x_i \in D$ for $i = 1, 2, ..., w$, we have*

$$\prod_{i=1}^{w} \frac{1}{\left([1-x_i]_q + [x_i]_q\right)^{n_i - l_i}} \sum_{k_1=m-1}^{n_1} \sum_{k_2=m-1}^{n_2} \cdots \sum_{k_w=m-1}^{n_w} \prod_{i=1}^{w} \frac{\binom{k_i}{m}}{\binom{n_i}{m}} B_{k_1,k_2,\cdots,k_w;n_1,n_2,\cdots,n_w}(x_1, x_2, \cdots, x_w; q)$$

$$= \sum_{k_1=0}^{m} \sum_{k_2=0}^{m} \cdots \sum_{k_w=0}^{m} q^{\sum_{i=1}^{w} \binom{k_i}{2}} \prod_{i=1}^{w} \binom{x_i}{k_i} [k_i]_q! S(l_i, k_i; q).$$

Proof. To prove this theorem, we let $F_q(t)$ be the generating function of the q-extension of the second kind Stirling numbers as follows:

$$F_q(t) := \frac{q^{-\binom{k}{2}}}{[k]_q!} \sum_{i=0}^{k} (-1)^{k-i} \binom{k}{i}_q q^{\binom{k-i}{2}} e^{[i]_q t} = \sum_{n=0}^{\infty} S(n,k;q) \frac{t^n}{n!}$$

From the above, we have

$$\begin{aligned} S(n,k;q) &= \frac{q^{-\binom{k}{2}}}{[k]_q!} \sum_{i=0}^{k} (-1)^i q^{\binom{i}{2}} \binom{k}{i}_q [k-i]_q^n \\ &= \frac{q^{-\binom{k}{2}}}{[k]_q!} \Delta_q^k 0^n \end{aligned} \tag{2.18}$$

where $[k]_q! = [k]_q [k-1]_q \cdots [2]_q [1]_q$. It is easy to see that

$$[x]_q^n = \sum_{k=0}^{n} q^{\binom{k}{2}} \binom{x}{k}_q [k]_q! S(n,k;q) \tag{2.19}$$

and in similar way that

$$\left(\prod_{i=1}^{w} [x_i]_q\right)^m = \sum_{k_1=0}^{m} \sum_{k_2=0}^{m} \cdots \sum_{k_w=0}^{m} q^{\sum_{i=1}^{w} \binom{k_i}{2}} \prod_{i=1}^{w} \binom{x_i}{k_i} [k_i]_q! S(l_i, k_i; q). \tag{2.20}$$

Then, we obtain the desired result from (2.19) and (2.20). □

3. INTERPOLATION FUNCTION OF MODIFIED q-BERNSTEIN POLYNOMIALS FOR FUNCTIONS OF SEVERAL VARIABLES

Definition 3. *Let $s \in \mathbb{C}$ and $x_i \neq 1$ with $i = 1, 2, ..., w$. We define interpolation function of the polynomials $B_{k_1,k_2,\cdots,k_w;n_1,n_2,\cdots,n_w}(x_1, x_2, \cdots, x_w; q)$ as*

$$D_q(s, k_1, k_2 \cdots k_w; x_1, x_2 \cdots x_w) = (-1)^{\sum_{i=1}^{w} k_i} \prod_{i=1}^{w} \frac{[x_i]^{k_i}}{k_i!} \left(\sum_{j=1}^{w} [1 - x_j] \right)^{-s}. \tag{3.1}$$

Remark 4. *By substituting $w = 1$ into (3.1), we get*

$$D_q(s, k_1) = (-1)^{k_1} \frac{[x_1]^{k_1}}{k_1!} ([1 - x_1])^{-s}$$

where $D_q(s, k_1)$ is introduced by Simsek and Acikgoz cf. [14].

By using (3.1), we have $D_q(s, k_1, k_2 \cdots k_w; x_1, x_2 \cdots x_w) \to D(s, k_1, k_2 \cdots k_w; x_1, x_2 \cdots x_w)$ as $q \to 1$. Thus one has

$$D(s, k_1, k_2 \cdots k_w; x_1, x_2 \cdots x_w) = (-1)^{\sum_{i=1}^{w} k_i} \prod_{i=1}^{w} \frac{x_i^{k_i}}{k_i!} \left(w - \sum_{j=1}^{w} x_j \right)^{-s}. \tag{3.2}$$

By substituting $x_j = 1$ with $i = 1, 2, ..., w$ into the above, we have $D(s, k_1, k_2 \cdots k_w; x_1, x_2 \cdots x_w) = \infty$.

We now evaluate the i-th s-derivative of $D(s, k_1, k_2 \cdots k_w; x_1, x_2 \cdots x_w)$ as follows:

$$\frac{\partial^i}{\partial s^i} D(s, k_1, k_2 \cdots k_w; x_1, x_2 \cdots x_w) = \log^i \left(\frac{1}{w - \sum_{j=1}^{w} x_j} \right) D(s, k_1, k_2 \cdots k_w; x_1, x_2 \cdots x_w) \tag{3.3}$$

where $x_j \neq 1$ with $i = 1, 2, ..., w$.

Remark 5. *By taking $w = 1$, $q \to 1$ into (3.2), we arrive at the relation which was proved by Simsek and Acikgoz* [1]*,*

$$\frac{\partial^i}{\partial s^i} D(s, k_1; x_1) = \log^i \left(\frac{1}{1 - x_1} \right) D(s, k_1; x_1).$$

REFERENCES

[1] Acikgoz, M., and Araci, S., On the generating function of the Bernstein polynomials, in proceedings of the 8th International conference of Numerical Analysis and Applied Mathematics (ICNAAM 2010), AIP, Rhodes, Greece, March 2010.

[2] Acikgoz, M., and Araci, S., New generating function of Bernstein type polynomials for two variables, in proceedings of the 8th International conference of Numerical Analysis and Applied Mathematics (ICNAAM 2010), AIP, Rhodes, Greece, March 2010.

[3] Acikgoz, M., and Araci, S., A study on the integral of the product of several type Bernstein polynomials, IST Transaction of Applied Mathematics Modelling and Simulation, 2010, vol. 1, no. 1(2), ISSN 1913-8342, pp. 10-14.

[4] Acikgoz, M., and Simsek, Y., On multiple interpolation functions of the Nörlund-type q-Euler polynomials, Abstr. Appl. Anal. 2009, Art. ID 382574, 14 pp.

[5] Buyukyazici, İ., and İbikli, E., Bernstein polynomials of two variable functions, Graduate School of Natural and Applied Sciences, Department of Mathematics, 1999, 49 pages, Ankara, Turkey.

[6] Buyukyazici, İ., and İbikli, E., The approximation properties of generalized Bernstein polynomials of two variables, Applied Math. and Comput. **156** (2004) 367-380.

[7] Kim, T., Jang, L.-C., and Yi, H., Note on the modified q-Bernstein polynomials, Discrete Dynamics in Nature and Society **2010 (**2010**),** Article ID 706483, 12 pages.

[8] Kim, T., Choi, J., and Kim, Y-H., On the k-dimensional generalization of q-Bernstein polynomials, Proceedings of the Jangjeon Mathematical Society, vol. 14, no. 2, pp. 199-207, 2011.

[9] Kim,T., A note on q-Bernstein polynomials, Russian Journal of Mathematical Physics **18** (2011), no. 1.

[10] Kim, T., Choi, J., and Kim, Y-H., q-Bernstein polynomials associated with q-Stirling numbers and Carlitz's q-Bernoulli numbers, Abstract and Applied Analysis, **2010** (2010), Article ID 150975, 11 pages.

[11] Oruc, H., and Phillips, G. M., A generalization of the Bernstein polynomials, Proceedings of the Edinburgh Mathematical society (1999) 42, 403-413.

[12] Ostrovska, S., On the q-Bernstein polynomials, Adv. Stud. Contemp. Math. 11 (2) (2005), 193-204.

[13] Phillips, G. M., A survey of results on the q-Bernstein polynomials, IMA Journal of Numerical Analysis Advance Access published online on June 23, (2009), 1-12, doi:10.1093/imanum/drn088.

[14] Simsek, Y., and Acikgoz, M., A new generating function of q-Bernstein-type polynomials and their interpolation function, Abstract and Applied Analysis, volume 2010, Article ID 769095, 12 pages, doi: 10.1155/2010/769095.01-313.

[15] Simsek, Y., Acikgoz, M., Bayad, A., and Lokesha, V., q-Frobenius-Euler polynomials related to the (q-) Bernstein type polynomials, in proceedings of the 8th International conference of Numerical Analysis and Applied Mathematics (ICNAAM 2010), AIP, Rhodes, Greece, March 2010.